

# THE PURDUE LANDSCAPE REPORT

## In This Issue

- [Purdue Extension and the Purdue Master Gardener Program: Resources for Home Gardeners](#)
- [Be on the Lookout for Defoliated Viburnums and Viburnum Leaf Beetle](#)
- [Dutch Elm Disease](#)

## Purdue Extension and the Purdue Master Gardener Program: Resources for Home Gardeners

(John Orick, [orick@purdue.edu](mailto:orick@purdue.edu)) & (Karen Mitchell, [mitcheka@purdue.edu](mailto:mitcheka@purdue.edu))

The Cooperative Extension Service is a national network of land-grant colleges, universities, and the United States Department of Agriculture serving communities across the country by providing research-based information and educational programming. Purdue Extension serves Indiana residents in the following program areas: Agriculture and Natural Resources, Health and Human Sciences, Community Development, and 4-H Youth Development. Purdue Extension offices are located in all 92 Indiana counties!



Photo 1: Purdue Master Gardeners in Tippecanoe County maintaining an educational demonstration garden. (Photo courtesy of Purdue Ag Communications)

One of Purdue Extension's signature programs is the Purdue Master Gardener Program. The purpose of the Purdue Master Gardener Program is to train volunteers to assist Purdue Extension Offices with consumer horticulture education needs in local communities. Purdue Master Gardeners complete horticultural training in a variety of topics from plant and soil science to fruit, vegetable, and ornamental gardening. The Purdue Master Gardener Basic Training is conducted by Purdue Extension Educators in participating Indiana Counties. To become

a Purdue Master Gardener Intern, participants must complete the application and screening requirements, pay the local fees, complete the Purdue Master Gardener Basic Training, and pass an exam over the material presented with a score of 70% or above. In order to become certified, interns are required to complete at least 35 hours of volunteer activity approved by their local Purdue Extension Educator (Master Gardener County Coordinator). Volunteer activities performed by Purdue Master Gardeners fulfill the program's purpose of educating Indiana residents in the area of home horticulture and gardening. Purdue Master Gardener volunteers can earn Advanced Master Gardener levels for their work. Purdue Master Gardeners are required to complete a minimum of 12 volunteer hours and 6 educational training hours per year to remain active in the program and continue their training in home horticulture. During 2018, 2,808 Purdue Master Gardener volunteers contributed 175,641 volunteer hours valued at over \$4.0 million!

Purdue Master Gardeners make a difference in the lives of others by sharing their gardening knowledge. Purdue Master Gardeners across the state have coordinated educational bus tours, held social functions to learn about gardening from each other, taught youth gardening classes, and conducted gardening seminars to educate the public. Some Purdue Extension offices offer education centers where Purdue Master Gardener volunteers are trained to assist Purdue Extension with home horticulture questions from local residents. The local Purdue Extension Educator, Agriculture and Natural Resources, supervises and provides ongoing training for Master Gardeners who staff the education centers. These valuable volunteers learn about working with the public and receive advanced training in diagnosis of home horticulture related plant problems. In 2018, the majority of questions answered at the Tippecanoe County Education Center were regarding trees at 42% of the total questions followed by lawn questions at 23%. The most common question of 2018 was "Why is my tree dying?" Although there were a variety of reasons for trees dying in Tippecanoe County, the most common cause for tree dieback was root girdling, which is ultimately caused by improper planting. With the 2019 season just beginning, we are seeing a similar trend with more questions regarding trees.

| 2018 Education Center Questions |                     |
|---------------------------------|---------------------|
| Question Topic                  | Percentage of Total |
| Trees                           | 42%                 |
| Lawn Care                       | 23%                 |
| Insect ID & Control             | 15%                 |
| Fruit                           | 9%                  |
| Weed ID & Control               | 8%                  |
| Other                           | 3%                  |



Photo 2: Purdue Master Gardener, Kathryn Reeves of Tippecanoe County, answers home gardening questions in the Education Center. (Photo by Karen Mitchell)

Indiana residents with gardening, lawn, and landscape questions are encouraged to contact their local Purdue Extension office (<https://extension.purdue.edu/about#counties>)! A trained Purdue Extension Educator can assist you by providing research-based nonbiased information from Purdue Extension experts. Likewise, the Purdue Extension Consumer Horticulture website offers extension publications and articles to assist the home gardener (<https://www.purdue.edu/hla/sites/yardandgarden/>). People interested in the Purdue Master Gardener Program are encouraged to contact their local Purdue Extension Office for more information about a Purdue Master Gardener Basic Training scheduled in a county near them ([www.hort.purdue.edu/mg](http://www.hort.purdue.edu/mg)).

Related Websites:

Purdue Extension: <https://extension.purdue.edu/>

Purdue Extension County offices:  
<https://extension.purdue.edu/about#counties>

Purdue Extension Consumer Horticulture:  
<https://www.purdue.edu/hla/sites/yardandgarden/>

Purdue Master Gardener Program: [www.hort.purdue.edu/mg](http://www.hort.purdue.edu/mg)

Purdue Master Gardener Basic Trainings:  
<https://www.purdue.edu/hla/sites/master-gardener/events/categories/2019-purdue-mg-basic-trainings/>

Purdue Plant Doctor Apps: <https://www.purdueplantdoctor.com/>

Purdue Plant and Pest Diagnostic Lab:  
<https://ag.purdue.edu/btny/ppdl/Pages/default.aspx>

## Be on the Lookout for Defoliated Viburnums and Viburnum Leaf Beetle

(Cliff Sadof, [csadof@purdue.edu](mailto:csadof@purdue.edu)) & (Elizabeth Barnes, [barne175@purdue.edu](mailto:barne175@purdue.edu))

With their attractive fruit, flowers, and fall color, viburnums are

popular shrubs in Indiana landscapes. Indeed their popularity extends into our native forests, where the fruit and leaves of these native plants are food for birds and other wildlife. Viburnums enjoy a natural resistance to many pests including Japanese beetles. The arrival of the viburnum leaf beetle, *Pyrrhalta viburni*, in the US threatens the health of viburnums. Although recently established in Illinois, Michigan, Wisconsin and Ohio, the viburnum leaf beetle has yet to be formally detected in Indiana. Containing this beetle before it becomes established in Indiana will save time, money, and natural resources. If you think you have seen viburnum leaf beetle or its damage report it via the 1-866-NO-EXOTIC phone number, the [GLEDN app](#), or on the [EDDMaps website](#).

### Viburnum Leaf Beetle Identification

Beetles winter as eggs on stems. Their characteristic small rough bumps can be easily detected during the dormant season.



Figure 1. Viburnum leaf beetles lay eggs on stems. (Photo by P. Weston, Cornell University, Bugwood.org)

Eggs hatch in May into green larvae that skeletonize leaves. You may have noticed holes in leaves or the larvae feeding on plants. After completing their development in mid-June, larvae will drop to the ground where they transform into the pupae that will eventually become adults.



Figure 2. Spiny larvae of viburnum leaf beetles chewing holes in leaves. (Photo by M. Zubrik, Forest Research Institute, Slovakia, Bugwood.org)

Golden colored adults will emerge in July to resume feeding on leaves. Feeding and eventually egg laying will continue until frost.



Figure 3. Adult beetles feeding on leaves (Photo by W. Cranshaw, Colorado State University, Bugwood.org)

## Viburnum Leaf Beetle Management

Arrowwood and cranberry bush viburnums are most susceptible to this pest. Doublefile and Koreanspice viburnums are resistant to this insect. Control this pest in dormant season by removing stems with obvious egg masses. In March prior to bud break, apply horticulture oil at the dormant rate (4%) to kill eggs glued to stems. When larvae are present on leaves 2% applications of insecticidal soap or spinosad (Captain Jack's Deadbug) will kill larvae. These two methods are preferred because they spare many of the beneficial insects that can attack and manage populations of the viburnum leaf beetle. Other products like bifenthrin and permethrin can be used to kill adults and larvae but care should be taken to avoid applying product when shrubs are in flower to avoid killing bees and other pollinators.

---

## Dutch Elm Disease

(Janna Beckerman, janna@purdue.edu)



Although most cities contain an Elm

Street, very few cities can claim having mature American elms on that street, or any other. The American elm was a premier street tree: Tolerant of compacted soil, fast growing, long-lived, and with a beautiful vase-shaped form. It's very popularity led to its downfall. The introduction of Dutch elm disease (DED) in the 1900's began devastating the elm population, which fell like dominos due to its overabundance in the urban and forest landscape. Although no number will be determined, the loss of hundreds of millions of elms is considered a conservative estimate, at best.

**The elm host.** Native Elm species vary in their susceptibility to DED. American elm (*Ulmus americana* L.) is highly susceptible, cedar elm (*U. crassifolia* Nutt.), rock elm (*U. thomasi* Sarg.), September elm (*U. serotina* Sarg.), slippery elm (*U. rubra* Muhl.), and winged elm (*U. alata* Michx.), range from susceptible to somewhat resistant. Although immunity does not appear to exist in native elm populations, select individuals or cultivars have been identified and propagated. European and Asian elms are less susceptible than American elm.

**Symptoms.** The first symptoms of Dutch elm disease are often overlooked due to the position of wilting leaves on branches in the upper canopy of the tree. Affected leaves usually turn yellow, then brown, and fall prematurely, although some leaves may remain attached. Upon losing its leaves, a branch quickly dies. Trees infected late in the summer or early fall usually survive only to rapidly succumb to the infection the following year. The rate of the infection varies, but an entire tree can die within a few weeks.



Very early symptoms of Dutch Elm Disease



Dutch Elm Disease



Brown streaks in the outer sapwood


DED-infected trees can be readily identified by brown streaks in the outer sapwood after the bark is removed. Keep in mind that elms produce both spring and summer wood, and that as the season progresses, previous vascular staining may be overlaid by unstained wood. It is important to note that elms suffer from other disease, insect and abiotic disorders; Conclusive diagnosis of DED requires laboratory testing.

### Causal Agent

Two fungi are responsible for this disease: *Ophiostoma ulmi* and *Ophiostoma novo-ulmi*. *O. novo-ulmi* is often referred to as a "newer, more virulent" version of the originally introduced pathogens. Both pathogens are established in the United States. Both species of *Ophiostoma* grow in the xylem (water-conducting tissue) of elms. Characteristic wilting results when the plant produces tyloses, gums and resins that block the spread of the fungus. These tyloses expand into the plants xylem, blocking water flow, and resulting in wilting. The fungus then spreads in the xylem, by growing yeast-like and also causing blockage of the xylem vessels. *Ophiostoma* produces a stalked spore structure within the beetle galleries (called a coremium), which is an aggregate of conidiophores forming a sticky head with millions of asexual spores (conidia). These sticky spores are vectored by the beetle when it moves within the gallery and emerges from April through June, and is the primary means the pathogen spreads from infected to healthy trees either. Beetles exit the trees to



feed on small branch crotches of other elms, providing an infection court for the fungus. Mature, adult females lay eggs in the wood of healthy, infected and dying elms. Larvae hatch and feed, forming tunnels from the main brood gallery into the surrounding tissue.



**Vector.** Two species of beetles vector

the pathogen: The European elm bark beetle (*Scolytus multistriatus*) (Fig. below) is the primary vector, and the native elm bark beetle (*Hylurgopinus rufipes*), to a lesser extent, spread the fungus from infected to healthy elms. The more common European elm bark beetles overwinter as larvae in recently killed elm wood. Overwintering adults of the native elm bark beetle emerge in the spring and tunnel into the inner bark of branches two to four inches in diameter to feed prior to breeding, when branches are abandoned to lay their eggs. Native bark beetle produce galleries that are bored perpendicularly to the grain of the wood, with developing larvae feeding in the direction of wood grain. In northern areas (Minnesota, Wisconsin, New England, and most of Canada), the native elm bark beetle is the predominant vector.

**Root Graft Transmission.** The pathogen also spreads via root grafts formed between adjacent or nearby elm trees (up to 50 feet away) when the root systems of adjacent elms fuse or graft. Should one tree become infected with *Ophiostoma*, the fungus can readily be transmitted to the vascular system of any other via the root-grafted tree. It has been estimated that more than 90 percent of urban elms killed in DED epidemics became infected through root grafts coupled with elm monocultures.

## Management

**Sanitation.** The simplicity of the concept of sanitation belies the great difficulty of its implementation. In order to be successful, DED sanitation programs must be applied without prejudice to the entire community. This includes every urban tree, whether situated in a park, cemetery, vacant lot, or hedgerow. DED-killed trees must be removed, stumps must be ground, and the wood must be promptly destroyed. Any and all wood from an infected tree should be burned, buried, or chipped. Wood chips from DED-infected trees pose no risk of disease transmission. DED-infected elms that are saved for firewood must be debarked, as logs with attached bark provide breeding sites for beetles.

**Root Graft Disruption.** Before disrupting root grafts, be sure to identify the location of possible buried utilities prior to digging. Disruption of root grafts between infected and healthy elms prevents transmission, thereby disrupting the spread of the disease. Determine the line of root graft disruption between adjacent trees by marking a line equidistant between the infected and adjacent tree. This line indicates where trenching or soil

fumigation placed should be done. Trenching can be performed with a trencher, ditch witch, vibratory plow, or a sharp spade, and should be dug to a depth of at least four feet.

In much of the urban landscape, trenching is not an option due to power lines, or concrete barriers. In this instance, chemical fumigants (soil sterilants) can be used to disrupt root grafts. One-inch diameter holes should be dug to a depth of two feet deep and placed every six to twelve inches between the two trees at a predetermined site equidistant to both trees. When concrete serves as a barrier, holes should be angled underneath the concrete to effectively distribute the fumigant. As always, follow label directions.

**Pruning.** Fresh pruning wounds attract beetles that may potentially vector the fungus that causes DED. Because of this, avoid unnecessary pruning whenever the beetles are active (early spring through late fall). If pruning is necessary, be sure to apply wound paint to prevent beetle/fungus access to wounds. This is one of the only instances when wound paint is warranted.



**Fungicide injection.** Although

fungicide injections have been successfully used for decades to protect and treat trees against DED, this option has not garnered wide-spread acceptance. First off, fungicide injections are more successful in preventing infection than treating already infected trees. Therapeutic treatment with either Arbortect, or Alamo is rarely successful when more than 25% of the canopy is symptomatic.

Fungicides are injected into the vascular system of elms in one of three ways: Microinjection forces a reduced volume of fungicide into holes drilled into the stem or base of the tree. Macroinjection is forceful injection of large volumes of a diluted fungicide or via infusion (without pressure) of large volumes of dilute chemical solutions into holes drilled in the stem or base of the tree. A third injection technique uses plastic capsules containing concentrated fungicide. Capsules are tapped into holes in the root flare.

To date, at least six chemicals are registered in the USA for injection to manage DED, and although it is legal for homeowners to apply injection treatments to their own trees, success rates are not as high.

**Adjuvants as Chemical Carriers.** The recent introduction, and subsequent labeling of Pentra-bark (Polyalkylene Modified Heptamethyltrisiloxane and nonionic surfactants) is changing how arborists can treat trees. Pentra-bark is an adjuvant that allows the transport of chemicals from the bark to the cambium of trees. To date, there are no data that demonstrate efficacy of this approach to manage Dutch elm disease.

## DED-Resistant Elms

Fungicides and sanitation have had limited success in the management of DED. Any long-term redeployment of elms in North America requires the identification and breeding of resistant elms. The use of European and Asian elms to introgress resistance, and the identification and propagation of DED-tolerant or resistant American has resulted in the identification of several cultivars showing high levels of DED tolerance and/or resistance.

In the decades since DED was introduced, American elms have been identified, and hybrid elms have been bred to be resistant to the disease, and these plants are now readily available to the public. It is important to note that resistance is not immunity, and resistance is impacted by climatic conditions (such as drought), bark beetle population, and surrounding sanitation, site, and overall tree health.

Link to [American Elm Clones of Importance in Dutch Elm Disease Tolerance Study](#) by Linda M. Haugen and Susan E. Bentz

Managing DED- resistance as a genetic resource requires careful

deployment of these varieties. Resistance is not the same as immunity, and stressed or poorly maintained trees will get infected. To minimize the likelihood of this, trees should be used more as specimens to diversify a landscape, as opposed to the monocultures that were once planted. Although these elm monocultures are visually pleasing, overuse of any cultivar or species should be avoided, to increase overall biodiversity and long-term ecosystem stability.

#### References:

Dunn, C.P. 2000. The Elms: Breeding, Conservation, and Disease Management. Kluwer Academic Publishers.

Stennes, M. and Haugen, L. 1999. Fungicide injections to control Dutch elm disease. Plant

Diagnostic Quarterly (PDQ) 20 (2):29-38

Sinclair, W. and Lyon, H. 2005. Diseases of Trees and Shrubs

It is the policy of the Purdue University that all persons have equal opportunity and access to its educational programs, services, activities, and facilities without regard to race, religion, color, sex, age, national origin or ancestry, marital status, parental status, sexual orientation, disability or status as a veteran. Purdue is an Affirmative Action Institution. This material may be available in alternative formats. 1-888-EXT-INFO Disclaimer: Reference to products in this publication is not intended to be an endorsement to the exclusion of others which may have similar uses. Any person using products listed in this publication assumes full responsibility for their use in accordance with current directions of the manufacturer.

Purdue Landscape Report © Purdue University - [www.purduelandscape.org](http://www.purduelandscape.org)

Editor: Kyle Daniel | Department of Horticulture and Landscape Architecture, 625 Agriculture Mall Dr., West Lafayette, IN 47907